

KANAZAWA-93-07

August, 1993

Differential Calculi on $\hbar$ -deformed Bosonic and Fermionic Quantum Planes

Tatsuo Kobayashi

Department of Physics, Kanazawa University,
Kanazawa, 920-11, Japan

Abstract

We study differential calculus on $\hbar$ -deformed bosonic and fermionic quantum space. It is shown that the fermionic quantum space involves a parafermionic variable as well as a classical fermionic one. Further we construct the classical $su(2)$ algebra on the fermionic quantum space and discuss a mapping between the classical $su(2)$ and the $\hbar$ -deformed $su(2)$ algebras.

1. Introduction

Quantum groups and quantum algebras [1-4] have attracted much attention in theoretical physics and mathematics, such as statistical models, integrable models, conformal field theories, knot theory and so on [5-8]. Quantum space was introduced to represent quantum groups [9]. Differential calculus on the quantum space was also studied [10-15]. It is intriguing also from the viewpoint of non-commutative geometry. The quantum differential calculus is closely related with q -oscillators, which have been applied to various fields [16-18]. Further, we can obtain quantum groups $Sp_q(2n)$ and $SO_q(2n)$ using the q -deformed phase space which is defined through the differential calculus on the quantum space for $SL_q(n)$ [19]. Furthermore, on the quantum space quantum deformed algebras have been constructed, e.g., q -deformed Lorentz, Poincaré, conformal and superconformal algebras [20-25]. These analyses imply that the quantum space is interesting as applications of quantum groups and useful to show some aspects of quantum groups.

Recently other than the conventional q -deformation and its multi-parametric extension, another deformation was discovered in [26-29] and is called h -deformation. Further, differential calculus on an h -deformed quantum bosonic space has been considered in [30, 31]. The purpose of this article is to investigate in detail bosonic and fermionic differential calculus on h -deformed quantum planes, extending the above analyses on the q -deformed differential calculus to the h -deformed case. The differential calculus on the bosonic quantum space is studied and a comment is obtained from the viewpoint of a constrained system. In addition, we discuss a differential calculus on an h -deformed fermionic space. Ref. [25] shows that q -deformed fermionic coordinates and derivatives represent $su_q(2)$ algebra, which is related with Drinfeld-Jimbo basis by some mapping. Following the approach, we investigate algebra which is constructed in terms of the h -deformed fermionic elements.

This paper is organized as follows. In section two we review on general non-commutative differential calculus including the quantum space. Then using a new solution of the Yang-Baxter equation, we derive h -deformed bosonic differential calculus. Further some comments on the h -deformed space are given. In section three, similarly we derive h -deformed differential calculus on the fermionic quantum space. It is shown that we can represent the classical Lie algebra $su(2)$, using their fermionic variables and derivatives. Also we obtain a mapping between the classical algebra

$su(2)$ and the h -deformed algebra $su_h(2)$. Further we discuss an h -deformed $SO(4)$ group. The last section is devoted to conclusion and discussion.

2. Differential calculus on h -deformed bosonic space

A quantum space is a non-commutative space representing the corresponding quantum group. Ref. [11] clarified general non-commutative differential calculus including the quantum space and its analysis was extended to superspaces in [15]. We set up commutation relations between coordinates x^i and derivatives ∂_{x^i} as follows,

$$x^i x^j = B^{ij}_{k\ell} x^k x^\ell, \quad \partial_{x^k} x^i = \delta_k^i + C^{ij}_{k\ell} x^\ell \partial_{x^j}. \quad (2.1)$$

These matrices should satisfied the following relations:

$$B^{ij}_{pr} B^{rk}_{tn} B^{pt}_{\ell m} = B^{jk}_{pr} B^{ip}_{\ell t} B^{tr}_{mn}, \quad (\delta_k^i \delta_\ell^j - B^{ij}_{k\ell})(\delta_m^k \delta_n^\ell + C^{k\ell}_{mn}) = 0. \quad (2.2)$$

The former equation is called Yang-Baxter equation. We consider the case where $B^{ij}_{k\ell}$ is propotional to $C^{ij}_{k\ell}$. In this case we can write commutation relations of derivatives by B -matrix as follows,

$$\partial_{x^\ell} \partial_{x^k} = B^{ij}_{k\ell} \partial_{x^j} \partial_{x^i}. \quad (2.3)$$

Recently a new solution of the Yang-Baxter equation was discovered in [27] as follows,

$$\hat{R} = \begin{pmatrix} 1 & -h' & h' & hh' \\ 0 & 0 & 1 & h \\ 0 & 1 & 0 & -h \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (2.4)$$

Also in Ref. [26] and [28], the same type of the $\hat{R}$ -matrix was discovered in the case where $h = h'$ and $h = h' = -1$, respectively. New type of a quantum group $SU_h(2)$ was studied in [26-28] and the corresponding deformed algebra $su_h(2)$ has been constructed in [29]. We identify $\hat{R}$ with B in order to obtain h -deformed differential calculus. The $\hat{R}$ -matrix has ± 1 as eigenvalues. This fact leads to a condition that we should identify C with $\hat{R}$, too. Namely we hereafter study the case where

$$B^{ij}_{k\ell} = C^{ij}_{k\ell} = \hat{R}^{ij}_{k\ell}. \quad (2.5)$$

We restrict ourselves to the case where $h = h'$. It is easy to extend the following analysis to the case with two independent parameters, h and h' . We substitute eq.(2.4) into (2.1) and (2.3), so that we obtain the following commutation relations;

$$\begin{aligned} [x^1, x^2] &= h(x^2)^2, \quad [\partial_{x^1}, \partial_{x^2}] = h(\partial_{x^1})^2, \\ [\partial_{x^2}, x^1] &= hx^1\partial_{x^1} + hx^2\partial_{x^2} + h^2x^2\partial_{x^1}, \\ [x^i\partial_{x^i}, x^i] &= 1 - hx^2\partial_{x^1}, \quad [\partial_{x^1}, x^2] = 0. \end{aligned} \quad (2.6)$$

The above algebra becomes classical in the limit, i.e., $h \rightarrow 0$.

We can introduce h-deformed quantum group T_j^i such that the elements transform the coordinates and the derivatives like $T_j^i x^j$. Covariance of the commutation relations (2.6) under the transformation requires the following commutation relation of T_j^i ;

$$\hat{R}^{ij} T_{k\ell}^k T_m^\ell = T_k^i T_\ell^j \hat{R}^{k\ell}_{mn}. \quad (2.7)$$

Next we consider conjugation consistent with (2.6). Suppose that h is pure imaginary, i.e., $\bar{h} = -h$. Then we obtain the consistent conjugation as follows,

$$\begin{aligned} \overline{x^1} &= x^1 + hx^2, \quad \overline{x^2} = x^2, \\ \overline{\partial_{x^1}} &= -\partial_{x^1}, \quad \overline{\partial_{x^2}} = -\partial_{x^2} - h\partial_{x^1}. \end{aligned} \quad (2.8)$$

Through the conjugation we can define real coordinates and momenta as follows,

$$\begin{aligned} \hat{x}^1 &= x^1 + \frac{h}{2}x^2, \quad \hat{x}^2 = x^2, \\ \hat{p}_1 &= -i\partial_{x^1}, \quad \hat{p}_2 = -i(\partial_{x^2} + \frac{h}{2}\partial_{x^1}). \end{aligned} \quad (2.9)$$

They satisfy the following commutation relations;

$$\begin{aligned} [\hat{x}^1, \hat{x}^2] &= h(\hat{x}^2)^2, \quad [\hat{p}_1, \hat{p}_2] = h(\hat{p}_1)^2, \\ [\hat{p}_j, \hat{x}^j] &= -i - h\hat{x}^2\hat{p}_1, \quad [\hat{p}_1, \hat{x}^2] = 0, \\ [\hat{p}_2, \hat{x}^1] &= h(-i + \hat{x}^1\hat{p}_1 + \hat{x}^2\hat{p}_2 - h\hat{x}^2\hat{p}_1). \end{aligned} \quad (2.10)$$

The similar phase space algebra as well as (2.6) has been obtained in [30].

These h-deformed coordinates and momenta can be represented in terms of classical ones $\tilde{x}$ and $\tilde{p}$. Suppose that $\hat{x}^2 = \tilde{x}^2$ and $\hat{p}_1 = \tilde{p}_1$, then we have

$$\hat{x}^1 = \tilde{x}^1(1 - ih\tilde{p}_1\tilde{x}^2) + ih(\tilde{x}^2)^2\tilde{p}_2, \quad \hat{p}_2 = \tilde{p}_2(1 - ih\tilde{p}_1\tilde{x}^2) + ih(\tilde{p}_1)^2\tilde{x}^2. \quad (2.11)$$

The deformed phase space algebra such as eq.(2.10) might remind us of Dirac brackets of a constrained system [32]. Actually we can find a “constraint” which we can make identically equal to zero through the above commutation relations (2.10), as follows,

$$f = 1 - 2ih\hat{p}_1\hat{x}^2. \quad (2.12)$$

The function f commutes with $\hat{x}^2$ and $\hat{p}_1$ and satisfies commutation relations with the other elements as follows,

$$[f, \hat{x}^1] = i\hat{x}^2 f, \quad [\hat{p}_2, f] = i\hat{p}_1 f. \quad (2.13)$$

These relations seems to be somewhat different from the Dirac bracket, whose right hand side vanishes completely. But we can make f equal to zero identically. The above fact seems to show that $SU_h(2)$ is a ‘group’ which transforms the “constraint” f and its relations (2.13) covariantly.

3. Differential calculus on h-deformed fermionic space

In this section we discuss differential calculus on an h-deformed quantum fermionic space. Commutation relations of h-deformed coordinates θ^α and derivatives ∂_α are obtained by replacing $\hat{R}$ in (2.1) and (2.3) in terms of $-\hat{R}$ as follows,

$$\begin{aligned} \theta^\alpha \theta^\beta &= -\hat{R}^{\alpha\beta}_{\mu\nu} \theta^\mu \theta^\nu, \quad \partial_\nu \partial_\mu = -\hat{R}^{\alpha\beta}_{\mu\nu} \partial_\beta \partial_\alpha, \\ \partial_\mu \theta^\alpha &= \delta_\mu^\alpha - \hat{R}^{\alpha\beta}_{\mu\nu} \theta^\nu \partial_\beta. \end{aligned} \quad (3.1)$$

These relations are written explicitly as follows,

$$\begin{aligned} (\theta^1)^2 &= h\theta^1\theta^2, \quad \{\theta^1, \theta^2\} = (\theta^2)^2 = 0, \\ (\partial_1)^2 &= \{\partial_1, \partial_2\} = 0, \quad (\partial_2)^2 = h\partial_1\partial_2, \\ \{\partial_\alpha, \theta^\alpha\} &= 1 + h\theta^2\partial_1, \quad \{\partial_1, \theta^2\} = 0, \\ \{\partial_2, \theta^1\} &= -h\theta^1\partial_1 - h\theta^2\partial_2 - h^2\theta^2\partial_1. \end{aligned} \quad (3.2)$$

Further eq.(3.2) leads to the following trilinear relations;

$$(\theta^1)^3 = (\partial_2)^3 = 0. \quad (3.3)$$

The coordinate θ^1 and the derivative ∂_2 are parafermionic [33], while its derivatives ∂_1 and its coordinates θ^2 are nilpotent.

As discussed in the previous section, we can introduce h-deformed quantum group elements T^α_β which transform θ^α into $\theta'^\alpha = T^\alpha_\beta \theta^\beta$. The elements satisfy the same commutation relation as (2.7). This h-deformed quantum group is interesting also from the aspect that it transforms fermionic and parafermionic variables, i.e., θ^1 and θ^2 . Further, we can define a determinant of T^α_β using the h-deformed fermionic space as follows,

$$\theta'^1 \theta'^2 = \det T \cdot \theta^1 \theta^2. \quad (3.4)$$

The definition leads to

$$\det T = T^1_1 T^2_2 - T^1_2 T^2_1 + h T^1_1 T^2_1. \quad (3.5)$$

Eq.(3.5) coincides with the definition of the determinant in [28].

Ref.[25] shows that q-deformed fermionic coordinates and derivatives can represent q-deformed quantum algebra, e.g., $su_q(2)$. Here we apply the similar analysis to the h-deformed case. In the similar way to [25], we define the following generators;

$$L_+ = \theta^2 \partial_1, \quad L_- = \theta^1 \partial_2. \quad (3.6)$$

Using their commutation relation, we introduce a Cartan generator as follows,

$$L_0 = [L_+, L_-] = \theta^2 \partial_2 - \theta^1 \partial_1. \quad (3.7)$$

In addition to (3.7), these generators satisfy the following relations;

$$[L_0, L_\pm] = \pm 2L_\pm. \quad (3.8)$$

Eqs. (3.7) and (3.8) are nothing but the classical $su(2)$ algebra up to the normalization factor, although the coordinates and the derivatives are deformed. The generators act on the coordinates and the derivatives as follows,

$$\begin{aligned} [L_0, \theta^\alpha] &= (-1)^\alpha \theta^\alpha, \quad [L_0, \partial_\alpha] = -(-1)^\alpha \partial_\alpha, \\ [L_+, \theta^1] &= \theta^2, \quad [L_+, \partial_2] = -\partial_1, \\ [L_+, \theta^2] &= [L_+, \partial_1] = 0, \\ [L_-, \theta^1] &= -h\theta^1 L_0 - h^1 \theta^2 L_+, \end{aligned} \quad (3.9)$$

$$\begin{aligned}
[L_-, \theta^2] &= \theta^1 + h\theta^2 L_0, \\
[L_-, \partial_1] &= -\partial_2 + h\partial_1 L_0 + h\partial_1 L_0, \\
[L_-, \partial_2] &= -h\partial_2 L_0 - 3h^2\partial_1 + 3h^2\partial_2 L_+.
\end{aligned}$$

The actions of L_+ and L_0 are never deformed, while those of L_- are deformed.

In Ref.[29] a h-deformed $su(2)$ algebra has been constructed as follows,

$$\begin{aligned}
[H, X] &= \frac{2\sinh(hX)}{h}, \\
[H, Y] &= -\{Y, \cosh(hX)\}, \\
[X, Y] &= H.
\end{aligned} \tag{3.10}$$

The h-deformed algebra $su_h(2)$ can be related with the classical algebra $su(2)$ through the following mapping;

$$\begin{aligned}
X &= \frac{\log \tilde{L}_+}{h}, \\
H &= (\tilde{L}_+ - (\tilde{L}_+)^{-1})\tilde{L}_0, \\
Y &= \frac{h}{2}(\tilde{L}_+ - (\tilde{L}_+)^{-1})(2\tilde{L}_+\tilde{L}_- - h\tilde{L}_0),
\end{aligned} \tag{3.11}$$

where $\tilde{L}_0 = L_0/2$ and $\tilde{L}_\pm = L_\pm/\sqrt{2}$.

In Ref.[19] Zumino derived quantum groups $Sp_q(2n)$ and $SO_q(2n)$ from q-deformed bosonic and fermionic phase space algebra for $SL_q(n)$. That was extended to the supersymmetric case in [15]. In the similar way, here we discuss h-deformation of $SO(4)$. At first, we define h-deformed gamma matrices γ^α as

$$\gamma^\alpha = \partial_\alpha, \quad \gamma^{5-\alpha} = \theta^\alpha. \tag{3.12}$$

They satisfy the following h-deformed Clifford algebra;

$$\gamma^\alpha \gamma^\beta + \tilde{B}^{\alpha\beta}_{\mu\nu} \gamma^\mu \gamma^\nu = \eta^{\alpha\beta}, \tag{3.13}$$

where $\eta^{\alpha\beta}$ is an $SO(4)$ metric. The matrix $\tilde{B}$ is composed of $\hat{R}$ -matrix (2.4) as follows,

$$\begin{aligned}
\tilde{B}^{\alpha'\beta'}_{\mu'\nu'} &= -\hat{R}^{\alpha\beta}_{\mu\nu}, \quad \tilde{B}^{\alpha\beta}_{\mu\nu} = -\hat{R}^{\nu\mu}_{\beta\alpha}, \\
\tilde{B}^{\alpha\beta'}_{\mu'\nu} &= -\hat{R}^{\beta\nu}_{\alpha\mu}, \quad \tilde{B}^{\alpha'\beta}_{\mu\nu'} = -(\hat{R}^{-1})^{\alpha\mu}_{\beta\nu},
\end{aligned} \tag{3.14}$$

where $\alpha, \beta, \mu, \nu = 1, 2$ and $\alpha' = 5 - \alpha$. Ref.[15] shows that $\tilde{B}$ constructed through the above procedure satisfy the Yang-Baxter equation, if $\hat{R}$ is the solution of the Yang-Baxter equation. We could introduce h-deformed $SO(4)$ group which transform the relation (3.13) covariantly. Instead of deriving explicitly $SO_h(4)$, we here introduce h-deformed $SO(4)$ quantum space X^i ($i = 1 \sim 4$) whose commutation relations are written as $X^i X^j = \tilde{B}^{ij}_{k\ell} X^k X^\ell$. We have explicitly

$$\begin{aligned} [X^1, X^2] &= h(X^1)^2, \quad [X^1, X^3] = 0, \\ [X^1, X^4] &= [X^2, X^3] = -hX^3X^1, \\ [X^2, X^4] &= hX^4X^1 + hX^3X^2 + h^2X^3X^1, \quad [X^3, X^4] = -h(X^3)^2. \end{aligned} \tag{3.15}$$

The algebra has a center element $C \equiv X^4X^1 + X^3X^2$. Differential calculus on the above h-deformed space X^i could be similarly obtained. Their commutation relations are covariant under the $SO_h(4)$ transformation, as said the above. Elements of $SO_h(4)$ satisfy the same relation as (2.7) except $\hat{R}$ -matrix replacing $\tilde{B}$ -matrix.

4. Conclusion

We have studied here the differential calculi on the h-deformed bosonic and fermionic spaces. We have constructed the classical $su(2)$ algebra on the h-deformed fermionic space. The algebra is related with the h-deformed algebra $su_h(2)$. It is shown that $SU_h(2)$ is a 'group' which transforms fermionic and parafermionic variables into each other. This fact is very interesting in applications to the parastatistics. Also $SO_h(4)$ was discussed. Supersymmetric extension is also intriguing. It is easy to introduce h-deformed oscillators in the similar way to the above h-deformed differential calculus. For example, we can define a sort of deformed oscillators by identifying the h-deformed coordinates and derivatives with deformed annihilation and creation operators, respectively. Their applications to various fields are very interesting.

Acknowledgement

The author would like to thank T. Uematsu for numerous valuable discussions and reading the manuscript. He also thanks to P. P. Kulish, R. Sasaki and H. Terao for helpful discussions.

References

- [1] V. G. Drinfeld, Sov. Math. Dokl. **32**(1985)254.
- [2] M. Jimbo, Lett. Math. Phys. **10**(1985)63.
- [3] L. D. Faddeev, N. Yu. Reshetikhin and L. A. Takhtajan, Algebra and Analysis, **1**(1987)178.
- [4] P. P. Kulish and N. Yu. Reshetikhin, J. Sov. Math. **23**(1983)2435.
- [5] H. -D. Doebner and J. -D. Hennig eds., *Quantum Groups*, Lecture Notes in Physics **370**(Springer Verlag, 1990).
- [6] T. Curtright, D. Fairlie and C. Zachos eds., *Quantum Groups*, Proceedings of the Argonne Workshop (World Scientific, 1990).
- [7] L. A. Takhtajan, Adv. Studies Pure Math. **19**(1989)435; Lecture Notes in Physics **370**(1989)3.
- [8] P. P. Kulish ed., *Quantum Groups*, Lecture Notes in Mathematics **1510**(Springer Verlag, 1992).
- [9] Yu. I. Manin, Commun. Math. Phys. **123**(1989)163.
- [10] S. L. Woronowicz, Commun. Math. Phys. **111**(1987)613; **122**(1989)125. Publ. RIMS, Kyoto Univ. **23**(1987)117.
- [11] J. Wess and B. Zumino, Nucl. Phys. B (Proc. Suppl.) **18B**(1990)302.
- [12] A. Schirmacher, J. Wess and B. Zumino, Z. Phys. C **49**(1991)317.
- [13] A. Schirmacher, Z. Phys. C **50**(1991)321.
- [14] U. Carow-Watamura, M. Schlieker and S. Watamura, Z. Phys. C **49**(1991)439.
- [15] T. Kobayashi and T. Uematsu, Z. Phys. C **56**(1992) 193.
- [16] A. J. Macfarlane, J. Phys. A **22**(1989)4581.
- [17] L. C. Biedenharn, J. Phys. A **22**(1989)L783.
- [18] M. Chaiahian and P. P Kulish, Phys. Lett. **B234**(1990)72.

- [19] B. Zumino, Mod. Phys. Lett. **A6**(1991)1225.
- [20] W. B. Schmidke, J. Wess and B. Zumino, Z. Phys. C **52**(1991)471.
- [21] O. Ogievetsky, W. B. Schmidke, J. Wess and B. Zumino, Lett. Math. Phys. **23**(1991)233.
- [22] O. Ogievetsky, W. B. Schmidke, J. Wess and B. Zumino, Commun. Math. Phys. **150**(1992)495.
- [23] T. Kobayashi and T. Uematsu, Z. Phys. C **58**(1993)559.
- [24] T. Kobayashi and T. Uematsu, Phys. Lett. **B306**(1993)27.
- [25] T. Kobayashi, “Quantum Deformed $su(m|n)$ Algebra and Superonformal Algebra on Quantum Superspace” Preprint KUCP-59; to be published in Z. Phys. C.
- [26] S. Zakrzewski, Lett. Math. Phys. **22**(1991)287.
- [27] H. Ewen, O. Ogievetsky and J. Wess, Lett. Math. Phys. **22**(1991)297.
- [28] E. E. Demidov, Yu. I. Manin, E. E. Mukhin and D. V. Zhdanovich, Prog. Theor. Phys. Suppl. **102**(1990)203.
- [29] Ch. Ohn, Lett. Math. Phys. **25**(1992)85.
- [30] V. Karimipour, “The Quantum de Rham Complexes Associatd with $SL_h(2)$ ” Preprint.
- [31] A. Aghamohammadi, “The 2-Parametric Extension of $\hbar$ deformation of $GL(2)$, and The Differential Calculus on Its Quantum Plane” Preprint.
- [32] P. Dirac, *Lectures on Quantum Mechanics* (Yeshiva University, NY, 1964).
- [33] Y. Ohnuki and S. Kamefuchi, *Quantum Field Theory and Parastatistics* (Springer Verlag, Berlin, 1982).